

TM



RODENT KILLER

# Professional

FOR USE BY REGISTERED  
PEST CONTROL  
OPERATORS ONLY

Reg No. / Nr.: L9290 Act / Wet 36 / 1947

## LIQUID CONCENTRATE

### A HIGHLY ACTIVE

A HIGHLY ACTIVE ANTICOAGULANT LIQUID BAIT CONCENTRATE FOR THE CONTROL OF THE NORWAY RAT (*RATTUS NORWEGICUS*), ROOF RAT (*RATTUS RATTUS*), THE HOUSE MOUSE (*MUS MUSCULUS*).

### 'N HOOGS AKTIEWE

'N HOOGS EFFEKTIEWE VLOEISTOFKONSTRAAT BLOEDSTOLWEERMIDDELOKAAS VIR DIE BEHEER VAN DIE NOORWEESE ROT (*RATTUS NORWEGICUS*), DAK ROT (*RATTUS RATTUS*), DIE HUISMUIS (*MUS MUSCULUS*).

ACTIVE INGREDIENT:  
Brodifacoum (Anticoagulant).....2,5 g/l

AKTIEWE BESTANDEEL:  
Brodifakoum (Bloedstolweermiddel) ...2,5 g/l

Registered and Formulated by / Geregistreer  
en Geformuleer deur:  
Innovative Pest Management  
Co. Reg. No./Mpy. Reg. Nr.: 2001/00587/07  
P.O. Box 3131, Kenmare, 1745

UN No. 3026


VERY TOXIC



UITERS GIFTIG



Date of Manufacture:  
Vervaardigingsdatum:  
Batch No.: / Lot Nr.:



6 009673 20017

# Rodex

# Professional

#### RODEX LIQUID CONCENTRATE

##### WARNINGS: HANDLE WITH EXTREME CARE

Prevent access to bait by children, domestic animals and unformed persons. Store under lock and key away from food and feedstuffs. Poisonous if swallowed and upon prolonged contact with skin. Toxic to fish and wildlife. In case of poisoning, call a doctor and make this label available to him.

##### PRECAUTIONS:

Avoid all contact by mouth. Avoid contamination of any edible commodity, feeds, food utensils and drinking water. Set bait stations where these will be inaccessible to children and domestic animals. Wear rubber gloves and wash hands after handling. Rinse empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the tank before destroying the container. Destroy empty container and do not re-use it for any other purpose. Do not eat, drink or smoke while handling the concentrate. Bury remains at end of treatment and burn or bury rodent bodies.

##### SYMPTOMS OF POISONING:

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints. Blood in urine and faeces.

##### FIRST AID TREATMENT:

If swallowed induce vomiting by tickling the back part of the throat. Repeat until vomit is clear. Take patient IMMEDIATELY to nearest physician.

##### NOTE TO PHYSICIAN:

Intravenous Vitamin K1, 50 -150 mg on first day, in divided doses, depending on prothrombin time. Keep under medical supervision until prothrombin times return to normal. In severe cases, two transfusions of fresh blood.

##### DIRECTIONS FOR USE:

USE ONLY AS DIRECTED FOR USE IN THE HOME, ON THE FARM AND INDUSTRIAL PREMISES. EFFECTIVE AGAINST WARFARIN RESISTANT RATS AND MICE.

1. LIQUID BAIT: Mix 1 part RODEX LIQUID CONCENTRATE with 160 parts clean water (5 ml to 800 ml water), place mixed liquid poison into liquid bait dispensers and place in areas of infestation, out of reach of children and animals.

All bait stations are to be inspected daily for the first week and weekly thereafter. Permanent Baiting should be continued in heavily infested areas using not only RODEX LIQUID CONCENTRATE, but also combining with a cereal type bait. This can be obtained by following the method below, or by using RODEX RAT AND MOUSE PELLETS or RODEX GRAIN BAIT.

2. MANUFACTURE OF SOLID BAITS, using cereal type carriers such as crushed wheat, crushed maize, maize meal and other suitable dry bait carriers. Mix 8 ml RODEX LIQUID CONCENTRATE with 12 ml clean water, then add to 1 kg of bait carrier. Mix thoroughly. Place 30 -50 g of bait into each bait station for rat control. Bait stations should be sited 5 meters apart. For mouse control, each bait station should contain 10 -20 g of bait and sited 2 meters apart. Place baits along walls, in rodent burrows and areas of infestation. Weekly inspections of bait stations is very important to achieve effective control. Death normally occurs in 4 -12 days.

##### WAARSKUWINGS: HANTEER UITERS VERSIGTIG

Hou buite bereik van kinders, troeteldiere en ongeligte persone. Bewaar agter slot en grendel weg van voedsel en voer. Giftig indien ingesluk of met langdurige velkontak. Giftig vir visse en wild. In geval van vergiftiging, ontbied 'n dokter en stel hierdie etiket tot sy beskikking.

##### VOORSORGMATREELS:

Vermi enige mondelinge inname. Voorkom besoedeling van eetbare produkte, voer, etgerei en drinkwater. Kies die lokaasplasingpunte sodat dit ontgevaarlik is vir kinders en troeteldiere. Dra rubber handskoene wanneer lokaas hanteer word en was hande na hantering. Spoel leë houers 3 keer uit met 'n volume water wat gelykstaande is aan minstens 10% van die houer. Voeg spoelwater by die inhoud van die tenk voordat die houer verniel of word. Vernietig leë houer en moet nooit vir enige ander doel hergebruik nie. Moet nie eet, drink of rook terwyl die lokaas hanteer word nie. Begrawe knaagdierkarkasse en enige oortollige lokaas na laaste behandeling.

##### SIMPTEME VAN VERGIFTIGING:

Bloeding van die tandvliese en neus, onderhuidse bloeding, bleekheid, bloeding in en rondom die knie -en elmboog gewigte. Bloed in urine en ontlasting.

##### WOODHULP BEHANDELING:

Indien ingesluk, help vomiting aan deur die agterkant van die keel te kliele. Harhaal tot vomiting helder is. Neem pasiënt ONMIDDELLIK na naaste geneesheer.

##### NOTA AAN GENEESHEER:

Vitamen K1 (Fibmenadioon) is die teenmiddel. Dien 50-150 mg intravenus toe, afhanklik van protrombientyd. Daaglikse behandeling mag nodig wees vir etlike weke. In ernstige gevalle, klein oortappings van vers bloed.

##### GEBRUIKSAANWYSINGS:

##### GEBRUIK SLEGS SOOS AANGEDUI.

VIR GEBRUIK IN WOONHUISE, OP PLASE EN INDUSTRIELE PERSELE. DOELTREFFEND TEEN WARFARINBESTANDE ROTTE EN -MUISE.

1. VLOEISTOF LOKAAS: Meng een deel RODEX LIQUID CONCENTRATE met 160 dele skoon water (5 ml op 800 ml water). Plaas aangemaakte vloeibare gif in lokaashouers en plaas dit buite bereik van kinders en huisdiere. Alle lokaashouers moet daaglik ondersoek word vir die eerste week en weekliks daarna. RODEX LIQUID CONCENTRATE behoort op 'n permanente basis in stand gehou te word. Waar erge infestasië voorkom, moet RODEX LIQUID CONCENTRATE in kombinasie met 'n "graan tipe" lokaas gebruik word. Gebruik onsevermelde metode vir vermenging van lokaas, of gebruik RODEX RAT AND MOUSE PELLETS of RODEX GRAIN BAIT.

2. VOORBEREIDING VAN SOLIEDE AAS: Gebruik graantipe draers soos gemaalde koring, gemaalde meel, mielie-meel en ander geskikte droë bestanddele.

Meng 8 ml RODEX LIQUID CONCENTRATE met 12 ml skoon water en voeg by 1 kg droë lokaasdraer. Meng deeglik. Plaas 30 -50 g lokaas in elke lokaashouer vir die beheer van rotte. Lokaashouers behoort 5 meter uit mekaar geplaas te word. Vir muisebeheer, plaas 10 -20 g lokaas in elke lokaashouer en plaas 2 meter uitmekaar uit. Houers kan langs mure, in knaagdiergate en in areas waar infestasië voorkom, geplaas word. Weeklikse inspeksies van lokaashouers is belangrik vir effektiwiteit. Vrektes kom gewoonlik na 4 -12 dae voor.

DEVELOPED BY  
**IPM**

1 Litre