



Reg No. / Nr.: L9289 Act / Wet 30 / 1947

RODEX KILLER LIQUID BAIT

A HIGHLY ACTIVE
anticoagulant liquid bait concentrate for
the control of the Norway Rat (*Rattus*
norvegicus), Roof Rat (*Rattus* *rattus*), the
House Mouse
(*Mus Musculus*).

ACTIVE INGREDIENT

Brodifacoum (Anticoagulant).....0.75 g/l

AKTIEWE BESTANDDEEL:

Brodifacoum (Bloedstolweermiddel) ..0.75 g/l

REGISTERED AND FORMULATED BY:
GEREGISTREER EN GEFORMULEER DEUR:
Innovative Pest Management

Co. Reg. No.
Mpy. Reg. Nr.: 2001 /00587 /07
P.O. Box 3131
Kenmare
1745
Tel: (011) 769 1300

UN No. 3026



VERY TOXIC
UITERS GIFTIG



200ml

WARNINGS: HANDLE WITH EXTREME CARE.
Prevent access to bait by children and domestic
animals. Store under lock and key away from
food and feedstuffs. Poisonous if swallowed and
upon prolonged contact with skin. Toxic to fish
and wildlife. In case of poisoning, call a doctor
and make this label available to him.

PRECAUTIONS:

Avoid all contact by mouth. Avoid contamination
of any edible commodity, feeds, food utensils and
drinking water. Set bait stations where these will
be inaccessible to children and domestic animals.
Wear rubber gloves and wash hands after
handling. Rinse empty container three times with
a volume of water equal to a minimum of 10% of
that of the container. Add the rinsings to the tank
before destroying the container. Destroy empty
container and do not re-use it for any other
purpose. Do not eat, drink or smoke while
handling the concentrate. Bury remains at end of
treatment and burn or bury rodent bodies.

SYMPTOMS OF POISONING:

Bleeding from the gums and nose, bleeding under
the skin, pallor, haemorrhage into and around
elbow and knee joints. Blood in urine and faeces.

FIRST AID TREATMENT:

If swallowed induce vomiting by tickling the back
part of the throat. Repeat until vomit is clear and
free from smell of poison. Take patient
IMMEDIATELY to nearest physician.

Date of Manufacture:

Vervaardigingsdatum:

Batch No.: / Lot Nr.:

NOTE TO PHYSICIAN:

Intravenous Vitamin K1, 50–150 mg on first day,
in divided doses, depending on prothrombin
time. Keep under medical supervision until
prothrombin times return to normal. In severe
cases small transfusions of fresh blood.

DIRECTIONS FOR USE:

USE ONLY AS DIRECTED FOR USE IN THE
HOME, ON THE FARM AND INDUSTRIAL
PREMISES, EFFECTIVE AGAINST WARFARIN
RESISTANT RATS AND MICE.

1. INITIAL BAITING FOR HEAVILY

INFESTED AREAS: Mix 1 part RODEX
LIQUID BAIT with 15 parts clean water (65 ml
to 1 litre water), place mixed liquid poison into
liquid bait dispensers and place in areas of
infestation, out of reach of children and animals.
All bait stations are to be inspected daily for the
first week and weekly thereafter. Permanent
baiting should be continued in heavily infested
areas using not only RODEX LIQUID BAIT in
combination with RODEX RAT AND MOUSE
PELLETS and RODEX GRAIN BAIT.

2. PERMANENT BAITING:

Mix 1 part
RODEX LIQUID BAIT with 50 parts clean
water (20 ml to 1 litre). Place liquid poison into
liquid bait dispensers and place in areas of
infestation, out of reach of children and animals.
All bait stations are to be inspected daily for the
first week and weekly thereafter.

Death normally occurs in 4–12 days.



6 009673 200032