



Avi-Carbo DP

Reg. No. / Nr. L1703 + G402 Act No./ Wet Nr. 36 of / van 1947

READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

GROUP

1A

INSECTICIDE

'n Nawerkende kontak- en maaggif vir die beheer van insekte op verskillende gewasse soos aangedui. Ook 'n kontakmiddel vir die beheer van rooimye, tampans en luise op hoenders en hokvoëls en in hoenderhuise. Beheer ook bosluise en vlooië op honde en katte.

A residual contact and stomach poison for the control of insects on various crops as listed. Also contact remedy for the control of red mites, tampans and, lice on poultry and cage birds and in poultry houses. Will also control ticks and fleas on dogs and cats.



WARNING

Hazard Statements:

May be harmful if swallowed

Suspected of causing cancer

Very toxic to aquatic life

Precautionary Statements:

Do not handle until all safety precautions have been read and understood

Avoid release into the environment

Active Ingredient (s) / Aktiewe Bestanddeel (ele)

Carbaryl (Carbamate) ... 50 g/kg ... Karbaril (Karbamaat)



Griffon Poison Information Centre: +27
82 446 8946

REGISTRATION HOLDER / REGISTRASIEHOUER

AVIMA (Pty.) Ltd.

Reg. No. 1961/001744/07

18 Aschenberg Street | Chamdor | Krugersdorp
+27 11 769 1300

**Content
Inhoud**

**Batch No. / Lot Nr and Date of Manufacture / Datum van
Vervaardiging:** See printed information on container.

UN No. / VN Nr. 2757



WARNINGS:

GHS Hazard Statements:

- May be harmful if swallowed.
- Suspected of causing cancer.
- Very toxic to aquatic life.
- Allow the following number of days between last application and harvest:

Tobacco:	14 days
Vines:	14 days
Cotton:	7 days
Lucerne:	1 day

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by conditions such as abnormal soil, climatic and storage conditions; compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as by the method, time and accuracy of application. The registration holder, furthermore, does not accept responsibility for damage to crops, vegetation, the environment, or harm to man or animal or for the lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

GHS Precautionary Statements:

- Do not handle until all safety precautions have been read and understood.
- Avoid release into the environment.
- Wear impervious rubber gloves and boots, protective clothing and chemical safety goggles.
- If exposed or concerned: Get medical attention.
- Collect spillage.
- Store locked up.
- Dispose of contents/container to suitable landfill in accordance with local regulations.
- Do not inhale dust.
- Wash contaminated clothing daily.
- Wash with soap and water after use.
- Avoid drift of the dust onto other crops, grazing, rivers, dams and areas not under treatment.
- Clean applicator before using it with other remedies and dispose of water where it will not contaminate crops, grazing, rivers and dams.
- Destroy empty container and never use for any other purpose.
- Prevent contamination of food, feedstuff, drinking water and eating utensils.

RELEVANT SUBSTANCES:

Inert	Weight %
Carbaryl	≥ 5

DIRECTIONS FOR USE: Use only as directed.

Red Mites and tampans in buildings: Dust all poultry houses thoroughly and pay particular attention to the floor, new litter, perches, nests, nest hay, cracks in wood, and walls.

Lice on birds: Dust under each wing and around vent.

Ticks and fleas on dogs & cats: Sprinkle powder lightly on the hair and rub in well. In the case of cats, the powder must be brushed out afterwards. Repeat weekly or as often as necessary.

Directions for application:

CROP	PEST	DOSAGE kg/ha	REMARKS
COTTON	Red bollworm, Spring bollworm, Stainers, Syagrus beetle, Jassids & Aphids	15 – 20 kg	Full cover application when pest is noticed.
LAWNS	Caterpillar, Crickets	20 kg (20 g/10 m ²)	Apply when damage is noticed. A damp sack left on the lawn during the night will reveal infestation of lawn caterpillar when removed in the morning. Light full cover application. Water in.
LUCERN	Caterpillar	15 – 20 kg	Full cover application when pest is noticed.
TOBACCO	American Bollworm, Slug	20 kg	Apply when pest is noticed. Direct application to flower head in the case of American bollworm.
VINES	Mealy bug	10 – 30 kg	Full cover application only to prevent build-up of mealy bug population.