

WARNINGS

- Handle with care.
- Harmful when swallowed or absorbed through the skin.
- Do not get in eyes, on skin or on clothing.
- Store in original container in a cool, dry place inaccessible to children and pets and away from food, feedstuffs and other chemicals.
- Keep containers tightly closed when not in use.
- Keep out of reach of children, uninformed persons and animals.
- In case of poisoning, call a doctor and present this label.

PRECAUTIONS

- Avoid contact with skin.
- Wear gloves when handling bait.
- After use and in case of accidental skin contact, wash thoroughly with soap and water.
- Do not expose children, pets or non-target animals to rodenticide.
- Store unused product out of reach of children and pets.
- Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtain bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock or other potentially destructive animals or in areas prone to vandalism.
- Do not contaminate ground water systems, surface water, ponds, waterways or ditches with rodenticide or its containers.
- Non-refillable container. Destroy empty container by perforation and burn and do not re-use for any other purpose.
- Collect unused, spoiled or unconsumed bait and dead rodents and dispose of in a responsible manner.

- Prevent contamination of food, feedstuffs, eating utensils and drinking water.

SYMPTOMS OF POISONING

Headache, confusion, personality change, tremors, convulsive seizures, respiratory distress.

FIRST AID TREATMENT

HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

Eyes – Flush with cool water for at least 15 minutes. If irritation develops, obtain medical assistance.

Skin – Wash with soap and water. If irritation develops, obtain medical assistance.

Ingestion – Call physician or emergency phone number immediately. Do not give anything by mouth or induce vomiting unless instructed by physician.

NOTE TO PHYSICIAN

If ingested, limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administrations of an osmotic diuretic and corticosteroid.

NOTE TO VETERINARIAN

Limit absorption by emesis or gastric gavage.

USE RESTRICTIONS

For control of Norway and roof rats and house mouse: Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. This rodenticide must be applied in tamper-resistant bait stations. Do not broadcast bait. This bait may be used in and around homes, agricultural, industrial, commercial, and public buildings.

Urban areas: This rodenticide may be used in and around the periphery of homes, industrial, commercial and public buildings. May also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. May also be used in alleys.

FASTRAC[®]

BLOX

A ready-for-use bait block for the control of roof and Norway rats and the house mouse in and around industrial, commercial and public buildings, in transport vehicles and in and around human dwellings and agricultural buildings.

Reg. No.: L9774 Act No. 36 of 1947

ACTIVE INGREDIENT:

BROMETHALIN (coumarin anticoagulant) 0.10 g/kg

NET MASS: 1.8 kg

Registered to:

MC Pharma (Pty) Ltd
Co Reg. No.: 1999/009355/07
P.O. Box 336
Wierda Park 0149
Pretoria, SOUTH AFRICA
Tel. No.: +27 (0) 12 668 3019

Poison Information Centre Emergency Tel. No. 0861 555 777

Manufactured by:



3699 Kinsman Blvd.
Madison, WI 53704 U.S.A.
www.thebelllabs.com
MADE IN USA

UN number: 3808.99.00



**CAUTION
INADHURU**

Non-urban areas: This rodenticide may only be used in and around homes and agricultural buildings.

DIRECTIONS FOR USE: Use only as directed

GENERAL:

Highest bait consumption is expected to occur on the first day or two after treatment begins. After several days, inspect the placement and replenish bait at sites where there is evidence of heavy feeding or where there is continued evidence of rodent activity. Normally, the volume of bait consumed will be significantly less with bromethalin baits than with anticoagulant baits. Although one night's feeding will usually provide a lethal dose, maintain bait for at least one week to provide all rodents a chance to feed on the bait. Continue baiting until all signs of feeding have ceased.

SELECTION OF TREATMENT AREAS:

Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain, direct sunlight and other extreme climatic conditions. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

MICE: Place 1 block per placement. Space placement at 2.5 to 3.5 meter intervals. Two blocks may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

RATS: Place 2 to 12 blocks (usually at intervals of 4.5 to 9.2 meters) per placement. Adjust the amount of bait applied to the level of rat feeding expected at each bait placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

FOR USE IN SEWER: Thread wire through block and securely attach to a stationary structure such as the bottom step of a manhole ladder or a sewer grate. Allow just enough wire for the block to rest on man-

hole benching. If benching is not present, suspend block a few centimeters above the high water mark of the ladder. Securing the block in this manner will prevent removal by rats or water. Use at least 15 blocks per manhole. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead, exposed animals and leftover bait. To prevent re-infestation, limit sources of rodent food, water and harbourage as far as possible. If re-infestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

Each bait block in this container weighs nearly 15 g.

STORAGE AND DISPOSAL

Do not contaminate water, food or feedstuffs by storage or disposal.

Storage: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide disposal:

In accordance with local and national regulations. Do not allow material to contaminate ground water system. Do not contaminate surface water.

Pesticide container: Do not re-use empty container. Do not contaminate ponds, waterways or ditches with chemical or used container. Destroy empty container by perforation and burn. If burning, stay out of smoke. Never use for any other purpose.

Fastrac[®] is a registered trademark of Bell Laboratories, Inc. U.S.A.

032816/10-15

Product Code: 993218



6508-1

