

THIS REMEDY CONTAINS AN ANTI-COAGULENT THAT MAY CAUSE INTERNAL BLEEDING IF CONSUMED.
THIS REMEDY MAY ONLY BE USED IN TAMPER-RESISTANT BAIT STATIONS AND ONLY AS DIRECTED ON THIS LABEL.
OBTAIN, READ AND FOLLOW ALL SAFETY INSTRUCTIONS BEFORE USE.



WARNING

HAZARD STATEMENTS:

- May cause damage to organs (blood) through prolonged or repeated exposure.

PRECAUTIONARY STATEMENTS:

- **Wear protective gloves/protective clothing/eye protection/face protection. The wearing of gloves while handling bait is essential.**
- **Obtain special instructions before use.**
- Wash hands thoroughly after handling.
- Do not eat, drink or smoke when using this product.
- Do not handle until all safety precautions have been read and understood.
- Specific treatment (see first aid section and read label).
- **If exposed or concerned, get medical advice.**
- Call a POISON CENTER/doctor if you feel unwell.
- **Store locked up.**
- **The use of tamper-resistant bait stations / boxes is a mandatory.**
- Avoid release to the environment.
- Dispose of contents/container to local and national regulations. Never reuse container. **Dispose of contents/container in a responsible manner to avoid accidental exposure of uninformed persons and animals.**
- Collect unused, spoiled or unconsumed bait and dead rodents and dispose of in a responsible manner.

DISCLAIMER

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as climatic and storage conditions; compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned, as well as by the method, time and accuracy of application.

The registration holder furthermore does not accept responsibility for damage to the environment, or harm to man or animal or for the lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration holder in the event of any uncertainty.

Consult the registration holder in the event of any uncertainty.

SYMPTOMS OF POISONING

Most important symptoms and effects, both acute and delayed. Ingestion of excessive quantities may cause nausea, vomiting, loss of appetite, extreme thirst, lethargy, diarrhoea, bleeding.

FIRST AID TREATMENT

Call a POISON CENTER/doctor if you feel unwell!

General information: When in doubt or if symptoms are observed, get medical advice.

Following inhalation: No special measures are necessary. Provide fresh air.

Following skin contact: Wash with soap and water.

Following eye contact: In case of eye irritation consult an ophthalmologist. Rinse immediately carefully and thoroughly with eye-bath or water.

Following ingestion: IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

IF SWALLOWED: immediately call a POISON CENTER!

Relevant Substance	Concentration (%)
Brodifacoum (coumarin)	
CAS N°	56073-10-0 C ≤ 0.005%



Reg. No. L8259 Act No. 36 of 1947 N-AR1499

For sale to and use only by registered pest control operators (PCO).

**READ THE LABEL BEFORE USE
AND**

KEEP OUT OF REACH OF CHILDREN AND ANIMALS
Effective against warfarin resistant Norway rats and house mouse.

POISON CENTER: +27 861 555 777

A ready-for-use bait block for the control of roof and Norway rats and the house mice in and around industrial, commercial and public buildings, in transport vehicles and in and around human dwellings and agricultural buildings.

Active Ingredient:

Brodifacoum (anti-coagulant) 0.05g/Kg (0.005%)

NETT WEIGHT: 8 kg

STORAGE AND DISPOSAL

Do not contaminate water, food or feedstuffs by storage or disposal.

Storage: Store only in original container in a cool, dry place inaccessible to children and pets. Keep container closed and away from other chemicals.

Pesticide disposal: In accordance with local and national regulations. Do not allow material to contaminate ground water system. Do not contaminate surface water.

Pesticide container: Do not re-use empty container. Do not contaminate ponds, waterways or ditches with chemical or used container. Destroy empty container by perforation and burn. If burning, stay out of smoke. Never use for any other purpose.

Collect unused, spoiled or unconsumed bait and dead rodents and dispose of in a responsible manner.

Registration Holder:

MC Pharma (Pty) Ltd. Co. Reg. No. 1999/009355/07
62 Constantia Avenue Mrandi, Centurion, Gauteng, 0157
Tel. No. +27 (0)12 668-3019 | www.mcpharma.co.za

Distributor:

Pathogen & Environmental Solutions (Pty) Ltd. Co. Reg. No. 2011/006448/07
Unit 37 Silver Park, Silver Street, Brackenfell Industria, Cape Town, 7560.
Tel. No. +27 (0) 21 982-8689 or +27 (0) 83 295 1313 or +27 83 272 3302 | info@pesafrica.net

Manufacturer:

Bell
LABORATORIES, INC.
www.belllabs.com
3699 Kinsman Blvd. Madison, WI 53704 U.S.A.
registration@belllabs.com | **MADE IN USA**

Date manufactured: see container

Expiry date: see container

Batch No.: see container

UN No.: not regulated

NOTE TO PHYSICIAN

Administer Vitamin K₁ intravenously, 50-150 mg depending on prothrombin time on the first day. In severe cases, small transfusions of fresh blood can be considered.

NOTE TO VETENARIAN

Administer Vitamin K₁ intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring prothrombin times.

Please note:

- Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations.
- These stations must be resistant to destruction by dogs and by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtain bait.
- If bait can be shaken from bait station when they are lifted, units must be secured or otherwise immobilized.
- Stronger bait stations are needed in areas open to hooded livestock or other potentially destructive animals or in areas prone to vandalism.
- Do not contaminate ground water systems, surface water, ponds, waterways or ditches with rodenticide or its container.
- Destroy empty container by perforation and burn and do not re-use for any other purpose.
- Collect unused, spoiled or unconsumed bait and dead rodents and dispose of in a responsible manner.
- Prevent contamination of food, feedstuffs, eating utensils and drinking water.
- Handle with care.

Urban areas: This rodenticide may be used in and around the periphery of homes, industrial, commercial and public buildings. May also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. May also be used in alleys.

Non-urban areas: This rodenticide may only be used in and around homes and agricultural buildings.

USE RESTRICTIONS:

For control of Norway and roof rats and house mice: Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

This rodenticide must be applied in tamper-resistant bait stations! Do not broadcast bait.

DIRECTIONS FOR USE - USE ONLY AS DIRECTED!

The wearing of gloves while handling bait is essential and the use of tamper-resistant bait stations / boxes is MANDATORY!

GENERAL:

Norway rats and the house mouse may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins.

SELECTION OF TREATMENT AREAS:

Determine areas where rats and mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain, direct sunlight and other extreme climatic conditions. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

MICE: Place 1 block per placement. Space placement at 2.5 to 3.5 metre intervals. Two blocks may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

RATS: Place 5 to 22 blocks (usually at intervals of 4.5 to 9.2 metres) per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rats' activity cease.

For use in sewer: Thread wire through block and securely attach to a stationary structure such as the bottom step of a manhole ladder or a sewer grate. Allow just enough wire for the block to rest on manhole benching. If benching is not present, suspend block a few centimetres above the high-water mark or secure block on a board supported by opposing steps of the ladder.

Securing the block in this manner will prevent removal by rats or water. Use at least 15 blocks per manhole. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

Follow up: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead, exposed animals and leftover bait. To prevent re-infestation, limit sources of rodent food, water and harbourage as far as possible. If re-infestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

Each bait block in this container weighs nearly 10g.



587AF-5

JAGUAR® is a registered trademark of Bell Laboratories, Inc. USA

062226/12-25 Product Code: 933325