

MARKED PROPERTY - NOT FOR UNAUTHORIZED REUSE. DO NOT REPLICATE.



MAXFORCE[®]

Prime

Insecticide

GROUP 4A INSECTICIDE

Reg. No. L8580 Act No. 36 of 1947

Before using this product read the label carefully

AN INSECTICIDAL GEL BAIT FOR THE CONTROL OF GERMAN COCKROACHES ONLY IN DOMESTIC PREMISES, PUBLIC HYGIENE AND FOOD STORAGE PREMISES. THE PRODUCT IS NON-VOLATILE AND NON-REPELLENT



Warning

ACTIVE INGREDIENTS:

Imidacloprid (chloro-nicotinyl).....21,5 g/kg

REGISTRATION HOLDER:

2022 Environmental Science ZA (Pty) Ltd. Reg. No. 2022/228577/07

P.O. Box 143, ISANDO 1600, SOUTH AFRICA

TEL: (011) 921 5911

MAXFORCE[®] is a registered trademark of the Bayer Group


UN No. 3077

10 x 30 g

envu™

Hazard statements: May cause an allergic skin reaction.
Very toxic to aquatic life with long lasting effects.

Precautionary statements: Wear protective gloves/
protective clothing/ eye protection/ face protection



Cockroach

ZA80918119G



The design and contents of this packaging are confidential and legally protected. Replication or any form of unauthorized copying is strictly forbidden. For permission to reproduce or distribute this packaging, written consent must be obtained from the registration holder.

MARKED PROPERTY - NOT FOR UNAUTHORIZED REUSE. DO NOT REPLICATE.



MAXFORCE®

Prime

30 g

Insecticide

GROUP 4A INSECTICIDE

Reg No. L8580

FOR FULL PARTICULARS SEE ENCLOSED PAMPHLET

Before using this product read the label carefully

AN INSECTICIDAL GEL BAIT FOR THE CONTROL OF GERMAN COCKROACHES ONLY IN DOMESTIC PREMISES, PUBLIC HYGIENE AND FOOD STORAGE PREMISES. THE PRODUCT IS NON-VOLATILE AND NON-REPELLENT



Warning

Hazard statements: May cause an allergic skin reaction. Very toxic to aquatic life with long lasting effects.

Precautionary statements: Wear protective gloves/ protective clothing/ eye protection/ face protection

ACTIVE INGREDIENTS: Imidacloprid (chloro-nicotinyl).....21,5 g/kg

REGISTRATION HOLDER:

2022 Environmental Science ZA (Pty) Ltd.

Reg. No. 2022/228577/07

P.O. Box 143, ISANDO 1600, SOUTH AFRICA

TEL: (011) 921 5911

MAXFORCE® is a registered trademark of the Bayer Group



6 005317 000983

UN No. 3077

ZA80928831H

The design and contents of this packaging are confidential and legally protected. Replication or any form of unauthorized copying is strictly forbidden. For permission to reproduce or distribute this packaging, written consent must be obtained from the registration holder.

MAXFORCE[®] Prime

Insecticide

GROUP 4A INSECTICIDE

Reg. No. L8580 Act No. 36 of 1947

Before using this product read the label carefully.

AN INSECTICIDAL GEL BAIT FOR THE CONTROL OF GERMAN COCKROACHES ONLY IN DOMESTIC PREMISES, PUBLIC HYGIENE AND FOOD STORAGE PREMISES. THE PRODUCT IS NON-VOLATILE AND NON-REPELLENT



Warning



Hazard statements

May cause an allergic skin reaction.

Very toxic to aquatic life with long lasting effects.

Precautionary statements

Wear protective gloves/ protective clothing/ eye protection/ face protection

ACTIVE INGREDIENTS:

Imidacloprid (chloro-nicotinyl).....21,5 g/kg

REGISTRATION HOLDER:

2022 Environmental Science ZA (Pty) Ltd. Reg. No. 2022/228577/07

P.O. Box 143, ISANDO 1600, SOUTH AFRICA

TEL: (011) 921 5911

MAXFORCE[®] is a registered trademark of the Bayer Group

DISCLAIMER:

ALTHOUGH THIS PRODUCT HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS, FAILURE THEREOF MAY ENSUE AS A RESULT OF A WIDE RANGE OF REASONS. IF THIS IS SUSPECTED, KINDLY NOTIFY THE REGISTRATION HOLDER.

WARNINGS:

Hazard statements

To avoid risks to human health and the environment, comply with the instructions for use

- Handle with care.
- Harmful when swallowed.
- May cause sensitization by skin contact. Harmful to fish and other animals.
- Store in a cool place and away from food and feedstuffs.
- Keep out of the reach of children, uninformed persons, and animals.
- The product will **NOT** control American Cockroaches effectively.
- Shelf-life: expires 2 years from date of manufacture.

PRECAUTIONS:

Precautionary statements

If skin irritation or rash occurs: Get medical advice/ attention.

- Avoid contact by mouth and skin.
- Wear rubber gloves during application.
- Wash with soap and water immediately after use or accidental skin contact.
- Do not eat, drink, or smoke whilst applying and before washing hands and face. Apply only in positions where it will be completely inaccessible to children and pets.
- Do not contaminate dams, drains, ponds, streams, and other waterways with the product or used containers. Do not apply gel to surfaces on which food or feed is stored, prepared, or eaten.
- Protect food preparing equipment and eating utensils from contamination during application. Destroy used container and do not re-use for any other purpose.

RELEVANT SUBSTANCES

Name	CAS-No. / EC-No. / REACH Reg. No.	Conc. [%]
Imidacloprid	138261-41-3	2,15
1,2-Benzisothiazol-3(2H)- one	2634-33-5 01-2120761540-60-xxxx	>= 0,05 - <= 0,1
reaction mass of 5-chloro-2-methyl-2H-isothiazol-3-on e and 2-methyl-2H-isothiazol-3- one (3:1)	55965-84-9	> 0,0015 - < 0,006

FIRST AID MEASURES:

General advice:	The nature of this product, when contained in commercial packs, makes spillage unlikely. However, if significant amounts are spilled then the following advice is applicable. Move out of dangerous area. If the patient is likely to become unconscious, place and transport in stable sideways position. Remove contaminated clothing immediately.
Skin contact:	Wash off immediately with plenty of soap and water, if available polyethyleneglycol 400, subsequently rinse with water. Take medical treatment if irritation develops and persist.
Eye contact:	Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Get medical attention if irritation develops and persists.
Ingestion:	Call a physician or poison control center immediately. Induce vomiting only, if: 1. patient is fully conscious, 2. medical aid is not readily available, 3. a significant amount (more than a mouthful) has been ingested and 4. time since ingestion is less than 1 hour. (Vomit should not get into the respiratory tract.) Call a physician or poison control center immediately.

NOTES TO PHYSICIAN:

Treatment:

- Treat symptomatically.
- Monitoring of respiratory and cardiac functions.
- In case of ingestion gastric lavage should be considered in cases of significant ingestions only within the first 2 hours.
- However, the application of activated charcoal and sodium sulphate is always advisable.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

- Suitable for use in commercial and industrial premises, food handling and processing establishments, restaurants, offices, store rooms, hospitals, schools, barracks, homes, hotels, ships, market places and warehouses.
- **Maxforce® Prime** contains a cockroach poison with a delayed action; therefore results are not immediately visible. An effect on the cockroach population can be expected within a week.
- Survey the area to be treated to identify centres of cockroach infestation.
- **Maxforce® Prime** is applied by spot-treatment in places in or around cockroach harbourages and where cockroaches are observed or hide.
- The gel must be applied in dark places and out of sight.
- **20 g:** Remove the cap of the cartridge tip. Apply pressure to the trigger to release droplets from the cartridge tip.
- **30 g:** **Maxforce® Prime** is dispensed using a purpose-built caulking gun, which is used as follows: Insert the cartridge into the gun frame. Remove the cap of the cartridge tip. Insert a dispensing tip as supplied with the caulking gun. A plastic dispensing tip should be used in and around electrical installations and equipment. Press the trigger of the caulking gun until the plunger contacts the piston of the cartridge. The next application of pressure to the trigger should release a droplet of gel from the dispensing tip.
- **Maxforce® Prime** can now be applied as required in and around buildings, equipment, furniture and electronic and mechanical equipment.

Application rates:

Bait application should equal approximately 0,25-0,5 g / spot and 10 to 20 spots / 10 m².

Infestation Level	Pest	Dosage Rate	Remarks
Low infestation	German Cockroach	0,25 g (3 mm diameter)	The lower application rate should be used for low and moderate German cockroach infestations or re- treatments.
High infestation	German Cockroach	0,5 g (6 mm diameter)	The higher rate should be used for more severe infestations of German cockroaches or first treatment.

Numerous smaller placements will provide faster control than a few larger spots.

- The gel will remain pliable and palatable to cockroaches as long as it is visible; usually up to 12 weeks.
- Where infestations are high inspect gel four weeks after treatment and replace if necessary. The life of the bait will be reduced under daylight conditions.

Maxforce® Prime can be integrated with space or surface spray treatments but must not be contaminated with other insecticides or placed on freshly sprayed surfaces.

For any product enquiry, contact 2022 Environmental Science ZA (Pty) Ltd. - 011 921 5911 (Office hours: 9am – 5pm).
For poisoning emergencies only, Contact Griffon Poison Information Centre: 082 446 8946.

MAXFORCE® is a registered trademark of the Bayer Group

© 2022 Environmental Science ZA (Pty) Ltd. All unauthorized reproduction of this label is forbidden.
www.za.envu.com

envu™
ZA809846181